

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

First synthesis of 4-substituted-benzenesulfonylcyanoguanidines

Masereel, Bernard; Lebrun, Philippe; Dogné, Jean-Michel; de Tullio, Pascal; Pirotte, Bernard; Pochet, Lionel; Diouf, Ousmane; Delarge, Jacques

Published in:
Tetrahedron Letters

Publication date:
1996

Document Version
Early version, also known as pre-print

[Link to publication](#)

Citation for pulished version (HARVARD):
Masereel, B, Lebrun, P, Dogné, J-M, de Tullio, P, Pirotte, B, Pochet, L, Diouf, O & Delarge, J 1996, 'First synthesis of 4-substituted-benzenesulfonylcyanoguanidines', *Tetrahedron Letters*, vol. 37, pp. 7253-7254.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



First Synthesis of 4-Substituted Benzenesulfonylcyanoguanidines

Bernard Masereel*, Philippe Lebrun[†], Jean Michel Dogné, Pascal de Tullio, Bernard Pirotte,
 Lionel Pochet, Ousman Diouf and Jacques Delarge


*Department of Medicinal Chemistry, University of Liège, 3, rue Fusch, 4000 Liège, Belgium

[†]Department of Pharmacology, University of Brussel, 808, route de Lennik, 1070 Brussels, Belgium

Abstract: N'-substituted-N-cyano-S-methylcarbamimidothioates react with 4-substitutedbenzene sulfonamides to give the corresponding sulfonylcyanoguanidines.

Copyright © 1996 Elsevier Science Ltd

The cyanoguanidine group is regarded as a non-classical bioisostere of the urea and thiourea function¹. This concept has been developed for the gastric antisecretory N'-alkyl-N-imidazolylalkylthioureas to give cimetidine². Conversely, N'-alkyl-N-substituted pyridyl-thioureas and -ureas were found to be as active as pinacidil, an antihypertensive cyanoguanidine^{3,4}. In the present study, we have applied this strategy to hypoglycemic sulfonylureas to obtain original drugs bearing a new moiety: the sulfonylcyanoguanidine function.




Scheme 1 Reagents : i, Y-NH₂ (1.5 equiv), ethanol, 25°C; ii, NaOH (1 equiv), DMF, dioxane, reflux.

The N''[4-(substitutedphenyl)sulfonyl]-N'-substituted-N''cyanoguanidines **5-20** (table 1) were prepared by refluxing an excess (1.2 equiv) of N'-substituted-N-cyano-S-methylcarbamimidothioate **2** with the sodium salt of 4-methylbenzenesulfonamide **35** or the 4-substitutedcarboxamidoethylbenzenesulfonamide **46** (scheme 1). The carbamidithioates **2** were synthesized by stirring a mixture of N-cyano-S,S'-dimethylimino carbonate **17** in ethanol with the required amine (Y-NH₂, 1.5 equiv).

The compounds **7**, **8**, **12** and **13** are the bioisosteres of tolbutamide, tolcyclamide, tolazamide and gliclazide, the main representatives of the hypoglycemic drugs of first chemical generation (X = CH₃)⁸. The antidiabetic sulfonylureas bearing a substituted carboxamidoethyl side-chain belong to the second chemical generation. Compound **16** is the bioisostere of glibenclamide, one of the most potent hypoglycemic drug⁸. The electron-withdrawing groups (SO₂ and =N-CN) should maintain the acidity of the moiety as compared to the sulfonylurea function. The ¹H-NMR spectra⁹ of **7** (δ = 5.6, 2H, br s) and **16** (δ = 4.7, 2H, br s) revealed that both protons of the cyanoguanidine moiety were not distinguishable and located with the residual water contained in DMSO-d₆. On the contrary, tolbutamide (δ = 10.41, 1H, br s; 6.30, 1H, br t) and glibenclamide (δ = 10.27, 1H, br s; 6.25 1H, br d), their sulfonylurea counterparts, clearly exhibited three different signals corresponding to the two urea protons and residual water. These data suggest tautomeric forms of the sulfonylcyanoguanidine moiety and a higher lability of their acidic proton as compared to their sulfonylurea bioisosteres. The chemical shift values lead to the proposal that **7** is probably more acidic than **16**, as it is

reported for tolbutamide ($pK_a = 5.3$)¹⁰ compared to glibenclamide ($pK_a = 6.8$)¹¹. All compounds were characterized by analytical and spectral methods⁹.

Table I : Synthesis of 4-substitutedbenzenesulfonylcyanoguanidines

				
N°	X	Y	Yield (%)	Mp (°C) ^a
5	CH ₃	C ₂ H ₅	17	131-133
6	CH ₃	(CH ₃) ₂ CH	72	132-134
7	CH ₃	CH ₃ (CH ₂) ₃	28	108-110
8	CH ₃	cyclohexyl	79	156-158
9	CH ₃	cycloheptyl	78	144-146
10	CH ₃	(±)4-CH ₃ cyclohexyl	52	163-165
11	CH ₃	(CH ₂) ₅ N	14	210-212
12	CH ₃	(CH ₂) ₆ N	22	193-195
13	CH ₃	1-azabicyclo[3,2,1]octane	19	186-188
14	C ₆ H ₅ CONHCH ₂ CH ₂	cyclohexyl	76	174-176
15	4-CH ₃ OC ₆ H ₄ CONHCH ₂ CH ₂	cyclohexyl	26	176-178
16	2-CH ₃ O,5-ClC ₆ H ₃ CONHCH ₂ CH ₂	cyclohexyl	50	92-95
17	2-CH ₃ O,5-ClC ₆ H ₃ CONHCH ₂ CH ₂	(±)4-CH ₃ cyclohexyl	64	85-87
18	2-furfurylCONHCH ₂ CH ₂	cyclohexyl	55	146-148
19	C ₆ H ₅ NHCONHCH ₂ CH ₂	cyclohexyl	37	108-110
20	4-ClC ₆ H ₄ NHCONHCH ₂ CH ₂	cyclohexyl	82	128-130

^a All compounds were crystallized from ethanol.

REFERENCES AND NOTES

- 1 Thornber, C.W. *Chem. Soc. Rev.* **1979**, 8, 563-580
- 2 Durant, G.J.; Emmett, J.C.; Gannellin C.R.; Miles, P.D.; Pearsons, M.E.; Prain, H.D.; White G.R. *J. Med. Chem.* **1977**, 20, 901-906
- 3 Manley, Q.W.; Quast, U. *J. Med. Chem.* **1992**, 35, 2327-2340
- 4 Takemoto, T.; Eda, M.; Okada, T.; Sakashita, H.; Matzno, S.; Gohda, M.; Ebisu, H.; Nakamura, M.; Fukaya, C.; Hihara, M.; Eikaku, M.; Yamanoushi, K.; Yokoyama, K. *J. Med. Chem.* **1994**, 37, 18-25
- 5 Arndt, F.; Martius, C. *Justus Liebigs Ann. Chem.* **1932**, 499, 228-287
- 6 Weber, H.; Aumüller, W.; Muth, K.; Weyer, R.; Heerdt, R.; Fauland, E.; Bänder, A.; Pfaff, W.; Schmidt, F.H.; Stork, H. *Drug Research / Arz. Forschung* **1969**, 19, 1326-1346
- 7 Hantzsch, A.; Wolvekamp, M. *Justus Liebigs Ann. Chem.* **1904**, 331, 264-297
- 8 Marchetti P.; Navalesi, R. *Clinical Pharmacokinetics* **1989**, 16, 100-128
- 9 The elemental analyses for C,H,N,S are within 0.4% of the theoretical values and were performed on a Carlo Erba 1108 analyzer. All new compounds gave IR and ¹H-NMR spectra in accordance with their proposed structure. 7: ¹H-NMR (80 MHz, d₆-DMSO) δ 7.65 (2H, d, J=8Hz), 7.30 (2H, d, J=8Hz), 5.60 (2H, m), 3.15 (2H, br s), 2.30 (3H, s), 1.65-1.25 (4H, m), 0.8 (3H, t). IR (KBr) 2199 cm⁻¹ (C≡N st). 16: ¹H-NMR (80 MHz, d₆-DMSO) δ 8.18 (1H, m), 7.75 (2H, d), 7.58 (1H, d), 7.4 (3H, dd), 7.05 (1H, d), 4.70 (2H, br s), 3.72 (3H, s), 3.25-3.65 (3H, m), 2.95 (2H, m), 1.80-0.95 (10H, m). IR (KBr) 2188 cm⁻¹ (C≡N st).
- 10 Häussler, A.; Hajdù, P. *Arch. Pharm.* **1958**, 291, 531-535
- 11 Hajdù, P.; Kohler, K.F.; Schmidt, F.M.; Spingler, H. *Drug Research / Arz. Forschung* **1969**, 19, 1381-1386

(Received in France 27 June 1996; accepted 20 August 1996)